

Investigation and Prevalence of Gastrointestinal Parasites of Equestrian Clubs Horses in Misurata, Libya

Layla O. Elmajdoub, Omar Mosaab, Om Assaad Alsaghir, and Sobhy Sorour Shimaa

ABSTRACT

Background and Objective: The horse has been a loyal friend and trusted partner of human beings. Horses are prone to infestation with both internal and external parasites. A cross-sectional study was performed to estimate the gastrointestinal parasite infecting horses in equestrian clubs.

Materials and Method: For the study, faecal samples were obtained from 50 randomly chosen horses of varying sexes and ages and analyzed using sedimentation techniques.

Results: The overall prevalence of egg/larva or cyst in the study area was 98.0% (49/50). Moreover, the mixed infection was higher than that of a single infection; there were significant differences between Protozoa and helminths ($P \leq 0.01$). The types of helminth parasites (egg/larva) detected were *Anoplocephala* spp. (17%), *Parascaris equorum* (17%), and *Moniezia* spp (20%) in increasing with the age of horses had a significant effect with ages above one year, moreover, there reported the oocysts of *Cryptosporidium parvum* (33%).

Conclusion: Hence, the present study's findings indicated a high prevalence of parasites compromising horses' health and welfare in the study area. Thus, proper screening and monitoring of the horses should be carried out regularly; regular and strategic anti-parasite programmers should be carried out.

Keywords: *Anoplocephala* spp, horses, intestinal parasites, helminths, protozoa, *Parascaris equorum*, and *Moniezia* spp.

Published Online: December 23, 2022

ISSN: 2684-5199

DOI : 10.24018/ejbio.2022.3.6.414

L. O. Elmajdoub*

Zoology Department, Faculty of Science,
Misurata University, Libya.

(e-mail:

elmajdoubayla@sci.misuratau.edu.ly)

O. Mosaab

Department of Veterinary Medicine,
College of Agriculture and Veterinary
Medicine, Qassim University, 51452
Qassim, Saudi Arabia.

O. A. Alsaghir

Zoology Department, Faculty of Science,
Misurata University, Libya.

S. S. Shimaa

Department of Parasitology, Faculty of
Veterinary Medicine, Kafrelsheikh
University, 33516, Egypt.

*Corresponding Author

I. INTRODUCTION

The digestive system of horses provides a target site for many intestinal parasite species; gastrointestinal parasites are responsible for severe pathological conditions, sometimes fatal, and reduced performance and physical illness, (Love *et al.*, 1999). Also provides a unique type of digestive system compared to other higher mammals. The stomach has four compartments. Among four compartments, the rumen is the largest part in the rumen partially chewed grass is stored and broken down into balls of cud, (Love *et al.*, 1999).

Horses has belonged to the equine group. It is found mainly in temperate, semi-arid or highland areas, Horses is herd animal and will happily live in groups with other animals of different species such as, sheep and goats. Horse is very friendly animals and enjoy the company of humans. Today its importance has scarcely diminished in parts of South America, Asia, Africa and Eastern Europe, (Addis *et al.*, 2017; Jajere *et al.*, 2016). Equids play crucial role in both urban and rural areas, providing agricultural energy and transport, and in many cases, the sole means of income generating for their resource-limited owners, (Getachew *et al.*, 2014).

Equines were infected with a wide range of intestinal parasites over 100 species, half of these species belonged to strongyles parasites (Lichtenfels *et al.*, 2008). In contrast,

while cestodes and trematodes are less occurring, *Eimeria* and *Isospora* species can also be encountered (Bowman, 2003). In the presence of a high infection rate of nematodes, chronicle catarrhal gastritis develops (Umue and Acici, 2009). Gastrointestinal parasites are one of the greatest limiting factors to successful horse rising throughout the world, (Jajere *et al.*, 2016). They are worldwide problem for both small and large-scale farmer with a greatest impact due to the availability of a wide range of agro-ecological factors suitable for diversified hosts and parasitic species, (Jajere *et al.*, 2016). The development and survival of helminths egg of larvae with faeces on pasture are depending on temperature and moisture, thus forming suitable environment for development of larvae of nematode and trematoda to infected stage, (AlAnazi and Alyousif, 2011).

Transmission of parasitic helminths is via the faecal-oral route, indirectly within contact. In addition, transmission can occur from person-to-person, from animal-to-person, animal-to-animal, moreover the diagnosis of gastrointestinal parasites through stool analysis is quite challenging even in severe infections. Moreover, the effects of gastrointestinal parasites are more evident in young and under nourished horses. Small numbers causes minimal damage, but large number pose a risk for colic and other symptoms. As a rule, older horses appear to develop immunity against the common gastrointestinal parasites and tend not to be affected by parasite

related problems as commonly as younger horses, (Jajere *et al.*, 2016).

Several studies have been conducted on gastrointestinal parasites from horses in many countries, such as Saudi Arabia in 20118, in Turkey (Negash *et al.*, 2021), in Western Australia (Boxell *et al.*, 2004), no gastrointestinal parasite fauna of horse in Libya is to date. According to estimates by the Libyan Ministry of Agriculture, Libya, the number of horses in Libya in 2016 exceeded 45,000 (Ministry of Agriculture, 2017). Parasitic helminths are more prevalent in foals, and young horses (Relf *et al.*, 2013), that is explained by an age acquired immunity. The males were shown to have higher infection compared to females (Buzatu *et al.*, 2017).

The present study was planned to diagnosis the gastrointestinal parasite fauna in some equines grazed in different Equestrian clubs in Misurata, Libya.

II. MATERIALS AND METHODS

A. Study Area

The present study was carried out from March to the end of November 2018 of some Equines from the Equestrian clubs in Misurata, Libya. Misurata is the third-largest city in Libya and occupies around 4000 km² of Libya's northwestern part (long. 15°6' E and lat. 32°23' N). The population in Misurata is estimated to be about 419,192 (Ministry of Agriculture, 2017).

B. Collection of Samples

A total of 50 faecal samples were collected and examined for equids from the Equestrian clubs for gastrointestinal helminth parasites (33 males and 17 from females). About 10 gram fresh faecal samples were collected directly from rectum using disposable polythene gloves, kept in plastic sachets. The necessary information was noted, such as faecal samples collection date, sex of horse, and preserved at 4 °C in the Zoology department laboratory.

C. Faecal Examination

The collected faecal samples were taken in clean Petri plates and thoroughly examined for colour, consistency, presence of blood, mucus, tapeworm segments and dead worms which sometimes, provided an important clue about the parasitic infection. The samples were examined microscopically using sedimentation method to detect parasitic infection (Solusby, 1982). One drop from the mixture was taken to prepare on the slide. The specimen was stained with Iodine wet mount solution and examined at 10X

and 40X objectives. In this way, two slides were designed from each sample were examined at 10X and 40X objectives of a microscope to detect eggs of helminths, protozoan's trophozoites or cysts of gastrointestinal parasites. For detection, the cryptosporidium cysts were made a smear on the slide and were air-dried; stained by modified Ziehl–Neelsen (Majewska *et al.*, 2004) examined using an oil immersion objective.

D. Data Analysis

Data were statistically analyzed using Pearson's Chi-square test with Yates continuity correction, performed by "R". In all cases 95% confidence interval (CI) and P<0.05 was considered for a statistically significant difference.

III. RESULTS

A. The Overall Prevalence of Gastrointestinal Parasites in Horses

From Table I, out of 50 samples, 49 (98%) were positive, 32 (97%) of males and 17 (100%) of females for the parasitic egg and cyst. The horse age was ranged from below one year to above ten years. The age ranged 1-4 years was the highest prevalence rate (36.7%) followed by 5-9 years (34.7%), based on statistical differences, was showed non-significant different between the prevalence rates according to sex and ages (P>0.05).

B. The Prevalence of Protozoan and Helminthic Parasites in Horses

Overall, horses were infected with protozoan and helminthic parasites. The mixed infection was the highest prevalence rate (63.3%) followed by an independent infection with protozoan parasites (36.7%), whereas, did not find any separate infection with helminths as illustrated in Table II. Based on horse sex, the mixed infection in female horses was higher than those in males (70.6% and 57.6%), respectively. In contrast, male horses' independent infection was higher than those in females (39.4% and 29.4%), respectively, as Table II. Statistical analysis found the highly significant difference between the protozoan and helminthic infection rates (P≤0.01). The highest prevalence rate with protozoan parasites at age range 5-9 years (36.6%) followed by aged ranged 1-4 years (34.4%). While the highest prevalence rate with helminth parasites at aged, 1-4 years (35.6%) followed by age 5-9 years (30.5%) with significant differences at (P≤0.01) as in Table III.

TABLE I: OVERALL PREVALENCE RATES OF GASTROINTESTINAL PARASITES IN HORSES

Ages	Examined Males	Examined Females	No. Male infected horses	No. infected female horses	Total infection rate
Below one year	6	0	6 (18.7%)	0 (0%)	6 (12.24%)
1-4 years	10	8	10 (31.3%)	8 (47.1%)	18 (36.7%)
5-9 years	10	7	10 (31.3%)	7 (41.2%)	17 (34.7%)
Above ten years	7	2	6 (18.7%)	2 (11.8%)	8 (16.3%)
Total	33	17	32 (97%)	17 (100%)	49 (98%)

TABLE IV: SIGNIFICANT DIFFERENCES WITH MEAN± S.E OF THE INTENSITY OF PROTOZOAN AND HELMINTHIC PARASITES IN HORSES

Intensity description	Protozoan infection			Helminthic infection		
	Rare	Moderate	Heavy	Rare	Moderate	Heavy
Mean±S.E	6.8±2.2**	3.67±1.7	5±3	3±0.71**	1.3±0.45**	0.82±0.44**

**Significant different (P≤ 0.01); S.E. Standard Error

TABLE II: OVERALL PREVALENCE RATES WITH MEAN± S.E OF PROTOZOAN AND HELMINTHIC PARASITES IN HORSES

	Protozoan infection rate	Helminthic infection rate	Mixed infection rate
Males	13 (39.4%)	0%	19 (57.6%)
Females	5 (39.4%)	0%	12 (70.6%)
Total	18 (36.7%)	0%	31(63.3%)
Mean±S.E	5.17±1.31	0%	1.7±0.35
Sign	**		**

**Significant different ($P \leq 0.01$); S.E. Standard Error

TABLE III: OVERALL PREVALENCE RATES OF PROTOZOAN AND HELMINTHIC PARASITES IN HORSES

Age	Protozoan infection rate	Helminthic infection rate
Below 1 year	8 (8.6%)	5 (8.5%)
1-4 years	32 (34.4%)	21 (35.6%)
5-9 years	34 (36.6%)	18 (30.5%)
Above 10 years	19 (20.4%)	14 (23.7%)
Sign		**

**Significant difference ($P \leq 0.01$)

C. The Intensity Infection rate of Protozoan and Helminthic Parasites in Horses

The intensity infection was ranged between rarely to heavy density with eggs and cysts, which showed the intensity infection with protozoa was higher than that with helminths. Table IV illustrated the significant differences ($P \leq 0.01$) between protozoa and helminths' intensity rates.

D. Overall Protozoan and Helminthic Parasites in Horses

Overall, horses were found to have been infected with protozoan parasites belonging to the three groups. The Sporozoa class (22.4 percent) saw the highest prevalence, and the Sarcodina class (7.32 percent) saw the lowest prevalence. The helminth parasites belonging to three groups, on the other hand, were isolated. The Nematode class was the highest prevalence rate (50 percent), and the Trematode class was the lowest prevalence rate (5.4 percent) in Table V. The statistical analysis found highly significant differences between the three protozoa classes ($P \leq 0.01$), whereas non-significant differences between the three classes of helminths ($P > 0.05$).

E. Overall Protozoan and Helminthic Species in Horses

Among Protozoa parasites, *Blantidium coli* showed the highest prevalence (34.4%) as Fig. 1. followed by *Cryptosporidium* sp. (33.3%), *Entamoeba coli* (16.1%), *Eimeria* sp. (13.9%), and the lowest prevalence rate *Isospora* spp (2.15%), (Table VI). The study showed an effect of sex on the prevalence of protozoan infection. In contrast, that indicates a higher prevalence in male horses. On the other hand, *Moniezia* sp. Showed the highest prevalence (20.4%) followed by *Parascaris equorum* as Fig. 2., *Anoplocephala* sp. *Trichostongylode* sp. (16.9%), and the lowest prevalence

showed by *Capillaria* sp. and *Gongulonema pulcurum* (3.4%). The study showed no effect of sex on the prevalence of helminth infection (Table VII). The statistical analysis showed the significant differences among the different species of protozoa and helminthic in both males and females ($P \leq 0.01$) in Fig. 3.




Fig. 1. Showing the trophozoite of *Blantidium coli*.Fig 2. Showing *Parascaris equorum* egg.Fig 3. Significant differences with Mean± S.E of protozoan and helminthic species in horses based on sex, showing the highly significant differences with *Blantidium coli* and *Cryptosporidium parvum* more than those other intestinal parasites infection.

TABLE V: PREVALENCE RATE AND SIGNIFICANT DIFFERENCES WITH MEAN± S.E OF CLASSES OF PROTOZOAN AND HELMINTHIC PARASITES IN HORSE

	Protozoan classes			Helminthic classes		
	Sporozoa	Sarcodina	Ciliophora	Nematoda	Cestoda	Trematoda
Prevalence rate	22.4%	7.32%	15.6%	50%	44.6%	5.4%
Mean±S.E	4.1±2.1	10.7±2.7**	5±2.52**	2.8±0.33	1.3±0.73	1±0.44 ^{NS}

**Significant different ($P \leq 0.01$) S.E. Standard Error

TABLE VII: PREVALENCE RATE OF HELMINTHIC SPECIES IN HORSES BASED ON SEX

	Total prevalence rate	Male prevalence rate	Female prevalence rate
<i>Parascaris equorum</i>	10 (16.9%)	4 (40%)	6 (60%)
<i>Anoplocephalasp</i>	10 (16.9%)	6 (60%)	4 (40%)
<i>Paranoplocephala mamillana</i>	3 (5.1%)	2 (66.7%)	1 (33.3%)
<i>Gongulonema pulcurum</i>	2 (3.4%)	1 (50%)	1 (50%)
<i>Trichostongyliode sp.</i>	10 (16.9%)	5(50%)	5(50%)
<i>Strongyliodes westri</i>	6 (10.2%)	4 (66.7%)	2 (33.3%)
<i>Paragonimus westermani</i>	3 (5.1%)	1 (33.3%)	2 (66.7%)
<i>Capllaira spp.</i>	2 (3.4%)	2 (100%)	0
<i>Moniezia sp.</i>	12 (20.4%)	7 (58.3%)	5 (41.7%)

TABLE VI: PREVALENCE RATE OF PROTOZOAN SPECIES IN HORSES BASED ON SEX

	Total prevalence rate	Male prevalence rate	Female prevalence rate
<i>Blantidium coli</i>	32 (34.4%)	22 (68.8%)	10 (31.3%)
<i>Entamoeba coli</i>	15 (16.1%)	11 (73.3%)	4 (26.7%)
<i>Cryptosporidium parvum</i>	31 (33.3%)	18 (58.1%)	13 (41.9%)
<i>Eimeria Spp.</i>	13 (13.9%)	10 (76.9%)	3 (23.1%)
<i>Isosopra spp.</i>	2 (2.15%)	1(50%)	1 (50%)

IV. CONCLUSION

In the conclusion, the current study's high parasitic prevalence rate could also be related to poor management or poor husbandry study area practices. In conclusion, this study confirmed that horses were found to be the most susceptible and infested by various gastro-parasites.

ACKNOWLEDGMENT

The authors are grateful to the Dean of the Zoology department, Misurata University, for providing the facilities for carrying out the research.

CONFLICT OF INTEREST

Authors declare that they do not have any conflict of interest.

REFERENCES

Addis, H., Gizaw, T.T., Minalu, B. A., & Tefera, Y. (2017). Cross sectional study on the prevalence of equine Strongyle infection Inmecha Woreda, Ethiopia. *International Journal of Advanced Research in Biological Sciences*, 2017 4(8): 68-77.

Adeppa, J., & K. J. Ananda. (2014). Incidence of gastro-intestinal parasites in horses of Shimoga region, Karnataka state. *J Parasit Dis*.

AL Anazi A.D. & Alyousif M.S. (2011). Prevalence of non-strongyle gastrointestinal parasites of horses in Riyadh region of Saudi Arabia, *Saudi Journal of Biological Sciences*, 2011, 18, 299-303.

Bowman D.D. (2003). *Georgis' Parasitology for Veterinarians*, 8nd edn, Elsevier Science, USA. pp: 215-216.

Boxell, A.C., Gibson, K.T., Hobbs, R.P., & Thompson, R.A.C. (2004). Occurrence of gastrointestinal parasites in horses in metropolitanPerth, Western Australia. *Aust. Vet. J.* 82 (1-2), 91-95.

Bucknell, D.G., Gasser, R.B., & Beveridge, I. (1995). The prevalence and epidemiology of gastrointestinal parasites of horses in Victoria, Australia. *International Journal for Parasitology*, 25(6): 711-724.

Buzato, M.C., Mitrea, I.L., Mironm L. & Ionita, M., (2015). Efficacy of two Anthelmintic Products on Strongyles in Horses from Stud Farms in Romania. *Agriculture and Agricultural Science Procedia*, 6, 293-298.

Buzatu M.C., Mitrea I.L., Gruianu A. & Ioniță, M. (2017). Investigating the Strongyle Populations, with Emphasis on Strongylus vulgaris (Nematoda: Strongylidae) in Romanian Horses, Based on Larval Cultures. *Bulletin UASVM Veterinary Medicinem*, 74(2), 186-192.

Chinwe, E.N., Amawulu, E., & Francis, G. C. (2019). Survey of Gastrointestinal Parasites and Ectoparasites of Horses (Equine Equine) in Port Harcourt and Abarka Polo Field, South Southern Nigeria. *International Journal of Science and Research*, 8(8),1471-1476.

Getachew, M., Alemayehu, F., Chala, C., Amare, B., Kassa, D., Burden, F., Wernery, R. & Werner, U. A. (2014). Cross sectional sero-survey of some infectious diseases of working equids in Central Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 6(9), 231-238.

Hasson, R.H. (2014). A study in the prevalence of horse "s helminthes parasitic infection in Baquba city, Digala. *International Journal of Recent Scientific Research*, 5(10), 1810-1813.

Mohammed Jajere, S., Rabana Lawal, J., Mohammed Bello, A., Wakil, Y., Aliyu Turaki, U., Waziri, I. (2016). Risk factors associated with the occurrence of gastrointestinal helminths among indigenous donkeys (Equus asinus) in Northeastern Nigeria. *Scientifica*.

Kornas, S., Marta, B., Vitalij, K. & Marta, S. (2010). Infection with large gastrointestinal parasites in slaughtered horses. *Bull Veterinary Institute polawy*, 54: 577-580.

Lichtenfels J.R., Kharchenko V.A., & Dvojnos G.M. (2008). Illustrated identification keys to strongylid parasites (Strongylidae: Nematoda) of horses, zebras and asses (Equidae). *Veterinary Parasitology*, 156, 4-161.

Love S., Murphy D., & Mellor D. (1999). Pathogenicity of cyathostome infection. *Veterinary Parasitology*, 85, 113-122.

Majewska AC, Solarczyk P, Tamang L, & Graczyk TK. (2004). Equine Cryptosporidium parvum infections in western Poland. *Parasitol Res*, 93(4), 274-8.

Matto, T.N., Bharkad, G.P. & Bhat, S.A. (2015). Prevalence of gastrointestinal helminth parasites of equids from organized farms of Mumbai and Pune. *J Parasit Dis*. 39 (2):179-185.

Mezgebu, T., Tafesi, K. & Tamiru, F. (2013). Prevalence of gastrointestinal parasites of horses and donkeys in and around Gondar town, Ethiopia. *Open Journal of Veterinary Medicine*, 2, 267-272.

Ministry of Agriculture. *Agriculture Statistical Year Book*, (2017), Libya.

Negash, W., Erdachew, Y. & Dubie T. (2021). Prevalence of Strongyle Infection and Associated Risk Factors in Horses and Donkeys in and around Mekelle City, Northern Part of Ethiopia. *Veterinary Medicine International*.

OLI, N. (2016). Prevalence of gastrointestinal parasites of horses (Equuscaballus Linnaeus, 1758) in seven vdc of rukum district, nepal. *Master thesis, Tribhuvan University Kirtipur, Kathmandu. Nepal*.

Ravi Prasad D., Janak R.S. & Kopila W. (2021). Prevalence of gastrointestinal parasites in equines of Mustang District, Nepal. *B I O D I V E R S I T A S*, 22(9), 3958-3963.

Relf, V.E., Morgan, E.R., Hodgkinson, J.E., & Matthews, J.B. (2013). Helminth egg excretionwith regard to age, gender and management *practices on UK thoroughbred studs.Parasitology*, 140, 641-652.

Solusby, E.J.L. (1982). *Helminths, Arthropods and Protozoa of Domesticated Animals . 7th ed. Bailliere Tindal. London*.

Tilahun, B., Nuraddis, I., Benti, D. & Tadele, T. (2014). Prevalence of helminth parasites of Horses in and around Hawasa town, southern Ethiopia. *College of agriculture and Vet. Med*, 5(1), 7-11.

Umar, Y.A., Maikaje, D.B., Garba, U.M. & Alhassan, M.A.F. (2013). Prevalence of gastrointestinal parasites in horses used for cadets training in Nigeria. *Journal of Veterinary Adv*, 3(2), 43-48.

- Umur, Ş. & Açııcı, M. A. (2009). Survey on helminth infections of equines in the Central Black Sea region, Turkey. *Turkish Journal of Veterinary and Animal Sciences*, 33(5), 373-378.
- Wannans, H.Y., Dawood, K.A. & Gassem, G.A. (2012). Prevalence of gastro- intestinal parasites of horses and donkeys in Diwaniyah. *Qudisiya Journal of Vet. Med. Sci.*, 11(1), 147-155.
- Yadav, K.S., Shukla, P.C., Gupta, D.V. & Mishra, A. (2014). Prevalence of gastrointestinal nematodes in horses of Jabalapur region. *Research Journal for Veterinary Practitioner*, 2(3), 44-48.